INDEX TO VOLUME 76

AUTHORS

Anderson, P. K., and R. M. R. Barclay:
Acoustic signals of solitary dugongs: physical characteristics and behavioral correlates, 1226
Antonelis, G. A.:
(see Ragen, T. J.)
Arends, A., F. J. Bonaccorso, and M. Genoud:
Basal rates of metabolism of nectarivorous bats (Phyllostomidae) from a semiarid thorn forest in Venezuela, 947
Arlettaz, R., G. Dändlinker, E. Kasybekov, J-M. Pillet, S. Rybin, and J. Zima:
Feeding habits of the long-eared desert bat, Otonycteris hemprichi (Chiroptera: Vespertilionidae), 873
Atramentowicz, M.:
Growth of pouch young in the bare-tailed woolly opossum, Caluromys philander, 1213
Baker, R. J.:
(see Jones, J. K., Jr.)
Barclay, R. M. R.:
(see Anderson, P. K.)
Barrett, G. W.:
(see Peles, J. D.)
Barrett, R. H.:
(see Beier, P.)
Barry, R. E., Jr.:
(see Bruseo, J. A.)
Bartlett, S. E.:
(see Perry, E. A.)
Batzli, G. O.:
(see Lin, Y. K.)
Beier, P., D. Choate, and R. H. Barrett:
Movement patterns of mountain lions during different behaviors, 1056
Bickham, J. W.:
(see Smolen, M. J.)
(see Morales, J. C.)
Bleich, V. C., and M. V. Price:
Aggressive behavior of Dipodomys stephensi, an endangered species, and Dipodomys agilis, a sympatric congener, 646
Boellstorff, D. E., and D. H. Owings:
Home range, population structure, and spatial organization of California ground squirrels, 551
Bolković, M. L., S. M. Caziani, and J. J. Protomastro:
Food habits of the three-banded armadillo (Xenarthra: Dasypodidae) in the dry Chaco, Argentina, 1199
Bonaccorso, F. J.:
(see Arends, A.)
Borrell, A.:
(see García, M.)
Bowers, M. A.:
Use of space and habitats of the eastern chipmunk, Tamias striatus, 12
Bowyer, R. T., J. W. Testa, and J. B. Faro:
Habitat selection and home ranges of river otters in a marine environment: effects of the Exxon Valdez oil spill, 1
Bozinovic, F.:
(see Sabat, P.)
Nutritional energetics and digestive responses of an herbivorous rodent (Octodon degus) to different levels of dietary fiber, 627
(see Lagos, V. O.)
Bradley, R. D.:
(see Jones, J. K., Jr.)
Bradshaw, S. D.:
(see Nagy, K. A.)
Braun, J. K., and M. A. Mares:
A new genus and species of phyllotine rodent (Rodentia: Muridae: Sigmodontinae: Phyllotini) from South America, 504
Britton-Davidian, J., J. Catalan, L. Granjon, and J-M. Duplantier:
Chromosomal phylogeny and evolution in the genus Mastomys (Mammalia, Rodentia), 248
Bronner, G. N.:
Cytogenetic properties of nine species of golden moles (Insectivora: Chrysochloridae), 957
Brooke, A. P.:
(see Studier, E. H.)
Brown, J. H.:
(see Valone, T. J.)
Bruseo, J. A., and R. E. Barry, Jr.:
Temporal activity of syntopic Peromyscus in the central Appalachians, 78
Bryant, P. J.:
Dating remains of gray whales from the eastern North Atlantic, 857
Burford, L. S.:
(see Lacki, M. J.)
Cameron, G. N.:
Temporal use of home range by the hispid cotton rat, 819
(see Randolph, J.C.)
Carr, S. M.:
(see Perry, E. A.)
Catlan, J.:
(see Britton-Davidian, J.)
Cavallini, P., and P. Serafini:
Winter diet of the small Indian mongoose, Herpestes auropunctatus, on an Adriatic island, 569
Caziani, S. M.:
(see Bolković, M. L.)
Chan, L-K.:
Extrinsic lingual musculature of two pangolins (Pholidota: Manidae), 472
Channell, R.:
(see Lomolino, M. V.)
Childs, J. E.:
Special feature: zoonoses, 663
(see Krebs, J. W.)
(see Mills, J. N.)
Childs, J. E., J. N. Mills, and G. E. Glass:
Rodent-borne hemorrhagic fever viruses: a special risk for mammalogists?, 664
Choate, D.:
(see Beier, P.)
Clements, C.:
(see Longland, W.S.)

INDEX TO VOLUME 76

AUTHORS

Anderson, P. K., and R. M. R. Barclay:
Acoustic signals of solitary dugongs: physical characteristics and behavioral correlates, 1226
Antonelis, G. A.:
(see Ragen, T. J.)
Arends, A., F. J. Bonaccorso, and M. Genoud:
Basal rates of metabolism of nectarivorous bats (Phyllostomidae) from a semiarid thorn forest in Venezuela, 947
Arlettaz, R., G. Dändlinker, E. Kasybekov, J-M. Pillet, S. Rybin, and J. Zima:
Feeding habits of the long-eared desert bat, Otonycteris hemprichi (Chiroptera: Vespertilionidae), 873
Atramentowicz, M.:
Growth of pouch young in the bare-tailed woolly opossum, Caluromys philander, 1213
Baker, R. J.:
(see Jones, J. K., Jr.)
Barclay, R. M. R.:
(see Anderson, P. K.)
Barrett, G. W.:
(see Peles, J. D.)
Barrett, R. H.:
(see Beier, P.)
Barry, R. E., Jr.:
(see Bruseo, J. A.)
Bartlett, S. E.:
(see Perry, E. A.)
Batzli, G. O.:
(see Lin, Y. K.)
Beier, P., D. Choate, and R. H. Barrett:
Movement patterns of mountain lions during different behaviors, 1056
Bickham, J. W.:
(see Smolen, M. J.)
(see Morales, J. C.)
Bleich, V. C., and M. V. Price:
Aggressive behavior of Dipodomys stephensi, an endangered species, and Dipodomys agilis, a sympatric congener, 646
Boellstorff, D. E., and D. H. Owings:
Home range, population structure, and spatial organization of California ground squirrels, 551
Bolković, M. L., S. M. Caziani, and J. J. Protomastro:
Food habits of the three-banded armadillo (Xenarthra: Dasypodidae) in the dry Chaco, Argentina, 1199
Bonaccorso, F. J.:
(see Arends, A.)
Borrell, A.:
(see García, M.)
Bowers, M. A.:
Use of space and habitats of the eastern chipmunk, Tamias striatus, 12
Bowyer, R. T., J. W. Testa, and J. B. Faro:
Habitat selection and home ranges of river otters in a marine environment: effects of the Exxon Valdez oil spill, 1
Bozinovic, F.:
(see Sabat, P.)
Nutritional energetics and digestive responses of an herbivorous rodent (Octodon degus) to different levels of dietary fiber, 627
(see Lagos, V. O.)
Bradley, R. D.:
(see Jones, J. K., Jr.)
Bradshaw, S. D.:
(see Nagy, K. A.)
Braun, J. K., and M. A. Mares:
A new genus and species of phyllotine rodent (Rodentia: Muridae: Sigmodontinae: Phyllotini) from South America, 504
Britton-Davidian, J., J. Catalan, L. Granjon, and J-M. Duplantier:
Chromosomal phylogeny and evolution in the genus Mastomys (Mammalia, Rodentia), 248
Bronner, G. N.:
Cytogenetic properties of nine species of golden moles (Insectivora: Chrysochloridae), 957
Brooke, A. P.:
(see Studier, E. H.)
Brown, J. H.:
(see Valone, T. J.)
Bruseo, J. A., and R. E. Barry, Jr.:
Temporal activity of syntopic Peromyscus in the central Appalachians, 78
Bryant, P. J.:
Dating remains of gray whales from the eastern North Atlantic, 857
Burford, L. S.:
(see Lacki, M. J.)
Cameron, G. N.:
Temporal use of home range by the hispid cotton rat, 819
(see Randolph, J.C.)
Carr, S. M.:
(see Perry, E. A.)
Catlan, J.:
(see Britton-Davidian, J.)
Cavallini, P., and P. Serafini:
Winter diet of the small Indian mongoose, Herpestes auropunctatus, on an Adriatic island, 569
Caziani, S. M.:
(see Bolković, M. L.)
Chan, L-K.:
Extrinsic lingual musculature of two pangolins (Pholidota: Manidae), 472
Channell, R.:
(see Lomolino, M. V.)
Childs, J. E.:
Special feature: zoonoses, 663
(see Krebs, J. W.)
(see Mills, J. N.)
Childs, J. E., J. N. Mills, and G. E. Glass:
Rodent-borne hemorrhagic fever viruses: a special risk for mammalogists?, 664
Choate, D.:
(see Beier, P.)
Clements, C.:
(see Longland, W.S.)

Contreras, L. C.:
(see Meserve, P. L.)
(see Lagos, V. O.)
Cook, J. A.:
(see Mouchaty, S.)
Cramer, K. L.:
(see Meserve, P. L.)
Cubbage, J.:
(see Tarpley, R. J.)
Cuthbert, F. J.:
(see Joshi, A. R.)
Dändliker, G.:
(see Arlettaz, R.)
Davidson, W. S.:
(see Perry, E. A.)
Davis, R. L., R. L. Lochmiller, and W. D. Warde:
Splenocyte subpopulations of weanling cotton rats (Sigmodon hispidus) are influenced by moderate protein intake, 912
DeLong, R. L.:
(see Stewart, B. S.)
Dickerman, A. W., and T. L. Yates:
Systematics of Oligoryzomys: protein-electrophoretic analysis, 172
Ding, H.:
(see Xu, Z.)
Dippenaar, N. J.:
Geographic variation in Mysorex longicaudatus (Soricidae) in the southern Cape Province, South Africa, 1071
Dumont, E. R.:
Enamel thickness and dietary adaptation among extant primates and chiropterans, 1127
Duplantier, J-M.:
(see Britton-Davidian, J.)
Duquette, L. S., and J. S. Millar:
Reproductive response of a tropical mouse, Peromyscus mexicanus, to changes in food availabilty, 596
Edge, W. D.:
(see Manning, T.)
Egozcue, J.:
(see García, M.)
Ellison, G. T. H.:
Thermoregulatory responses of cold-acclimated fat mice (Steatomys pratensis), 240
Engle, D. M.:
(see Lochmiller, R. L.)
Ernest, K. A.:
(see Mares, M. A.)
Estes, J. A.:
(see McShane, L. J.)
Etnier, S. A.:
(see Thewissen, J. G. M.)
Faro, J. B.:
(see Bowyer, R. T.)
Fayrer-Hosken, R. A.:
(see White, L. M.)
Fenn, M. G. P., and D. W. MacDonald:
Use of middens by red foxes: risk reverses rhythms of rats, 130
Fleck, D. W., and J. D. Harder:
Biology of marsupials in two Amazonian rain forests in northeastern Peru, 809
Gage, K. L., R. S. Ostfeld, and J. G. Olson:
Nonviral vector-borne zoonoses associated with mammals in the United States, 695
García, M., A. Borrell, M. Mudry, J. Egozcue, and M. Ponsá:
Prometaphase karyotype and restriction-enzyme banding in squirrel monkeys, Saimiri boliviensis boliviensis (Primates: Platyrrhini), 497
Gates, C.:
(see van Zyll de Jong, C. G.)
Gehring, J. A.:
(see Long, C. A.)
Genoud, M.:
(see Arends, A.)
George, J. C.:
(see Tarpley, R. J.)
Glass, G. E.:
(see Childs, J. E.)
Goodwin, H. T.:
Pliocene-Pleistocene biogeographic history of prairie dogs, Genus Cynomys (Sciuridae), 100
Gottelli, D.:
(see Sillero-Zubiri, C.)
Goyal, S. P.:
(see Pendharkar, A. P.)
Granjon, L.:
(see Britton-Davidian, J.)
Grant, W. E.:
(see Killion, M. J.)
Gutiérrez, J. R.:
(see Meserve, P. L.)
Hafner, D. J., and R. M. Sullivan:
Historical and ecological biogeography of Nearctic pikas (Lagomorpha:Ochotonidae), 302
Halle, S.:
Effect of extrinsic factors on activity of root voles, Microtus oeconomus, 88
Hammond, K. A., and B. A. Wunder:
Effect of cold temperatures on the morphology of gastrointestinal tracts of two microtine rodents, 232
Hansson, L.:
Size dimorphism in microtine populations: characteristics of growth and selection against large-sized individuals, 867
Harder, J. D.:
(see Fleck, D. W.)
Heinrich, R. E., and K. D. Rose:
Partial skeleton of the primitive carnivoran Miacis petilus from the early Eocene of Wyoming, 148
Helle, E., and K. Kauhala:
Reproduction in the raccoon dog in Finland, 1036
Hellgren, E. C.:
(see Ilse, L. M.)
Herman, T. B.:
(see Teferi, T.)
Herrera, S.:
(see Meserve, P. L.)
Heske, E. J.:
Mammalian abundances on forest-farm edges versus forest interiors in southern Illinois: is there an edge effect?, 562
Heske, E. J., G. I. Shenbrot, and K. A. Rogovin:
Spatial organization of Stylodipus telum (Dipodidae, Rodentia) in Dagestan, Russia, 800

Hickling, G. J.:
(see Moses, R. A.)
Hohn, A. A.:
(see Tolley, K. A.)
Hollander, R. R.:
(see Willig, M. R.)
Hosack, D. A.:
(see White, L. M.)
Hudson, M. K.:
(see Sabol, B. M.)
Hughes, J. J., D. Ward, and M. R. Perrin:
Effects of substrate on foraging decisions by a Namib Desert gerbil, 638
Huot, J.:
(see Samson, C.)
Ilse, L. M., and E. C. Hellgren:
Resource partioning in sympatric populations of collared peccaries and feral hogs in southern Texas, 784
Spatial use and group dynamics of sympatric collared peccaries and feral hogs in southern Texas, 993
Irvine, A. B.:
(see Tolley, K. A.)
Jacobi, C. L.:
(see Valone, T. J.)
Jaksic, F. M.:
(see Meserve, P. L.)
Jarrell, G. H.:
(see Tarpley, R. J.)
Johansson, A., O. Liberg, and L. K. Wahlström:
Temporal and physical characteristics of scraping and rubbing in roe deer (Capreolus capreolus), 123
Johnson, M. K.:
(see Schultz, S. R.)
Jones, J. K., Jr., R. D. Bradley, and R. J. Baker:
Hybrid pocket gophers and some thoughts on the relationship of natural hybrids to the rules of nomenclature and the Endangered Species Act, 43
Joshi, A. R., J. L. D. Smith, and F. J. Cuthbert:
Influence of food distribution and predation pressure on spacing behavior in palm civets, 1205
Kasybekov, E.:
(see Arlettaz, R.)
Kaufman, D. M.:
Diversity of New World mammals: universality of the latitudinal gradients of species and bauplans, 322
Kauhala, K.:
(see Helle, E.)
Kieser, J. A.:
Gnathomandibular morphology and character displacement in the bat-eared fox, 542
Killion, M. J., W. E. Grant, and S. B. Vinson:
Response of Baiomys taylori to changes in density of imported fire ants, 141
Kiyota, M.:
(see Ragen, T. J.)
Krajewski, C.:
(see Painter, J.)
Krebs, J. W., M. L. Wilson, and J. E. Childs:
Rabies—epidemiology, prevention, and future research, 681
Ksiazek, T. G.:
(see Mills, J. N.)
Kunz, T. H.:
(see Studier, E. H.)
Kunz, T. H., and S. K. Robson:
Postnatal growth and development in the Mexican free-tailed bat (Tadarida brasiliensis mexicana): birth size, growth rates, and age estimation, 769
Kvam, T.:
(see Strand, O.)
Lacki, M. J., L. S. Burford and J. O. Whitaker, Jr.:
Food habits of gray bats in Kentucky, 1256
Lagos, V. O.:
(see Meserve, P. L.)
Lagos, V. O., F. Bozinovic, and L. C. Contreras:
Micro-habit use by a small diurnal rodent (Octodon degus) in a semiarid environment: thermoregulatory constraints or predation risk?, 900
Lambertsen, R., N. Ulrich, and J. Straley:
Frontomandiblular stay of Balaenopteridae: a mechanism for momentum recapture during feeding, 877
Lang, B. K.:
(see Meserve, P. L.)
Leslie, D. M., Jr.:
(see Lochmiller, R. L.)
Lewis, S. E.:
Roost fidelity of bats: a review, 481
Lidberg, O.:
(see Johansson, A.)
Lima, M., and E. Páez:
Growth and reproductive patterns in the South American fur seal, 1249
Lin, Y. K., and G. O. Batzli:
Predation on voles: an experimental approach, 1003
Lochmiller, R. L.:
(see Davis, R. L.)
Lochmiller, R. L., D. G. Peitz, D. M. Leslie, Jr., and D. M. Engle:
Habitat-induced changes in essential amino-acid nutrition in populations of eastern cottontails, 1164
Löfgren, O.:
Niche expansion and increased maturation rate of Clethrionomys glareolus in the absence of competitors, 1100
Lomolino, M. V., and R. Channell:
Splendid isolation: patterns of geographic range collapse in endangered mammals, 335
Long, C. A., and J. A. Gehring:
Valais-goat color pattern in a masked shrew (Sorex cinereus) and lack of size dependence in pigmentation patterns, 937
Longland, W. S., and C. Clements:
Use of fluorescent pigments in studies of seed caching by rodents, 1260
Lucas, M. F.:
(see Peles, J. D.)
MacDonald, D. W.:
(see Fenn, M. G. P.)
Manning, T., W. D. Edge, and J. O. Wolff:
Evaluating population-size estimators: an empirical approach, 1149

Mares, M. A.:
(see Braun, J. K.)
Mares, M. A., and K. A. Ernest:
Population and community ecology of small mammals in a gallery forest of central Brazil, 750
McClure, P. A.:
(see Randolph, J.C.)
McNab, B. K.:
Energy expenditure and conservation in frugivorous and mixed-diet carnivorans, 206
McShane, L. J., J. A. Estes, M. L. Riedman, and M. M. Staedler:
Repertoire, structure, and individual variation of vocalizations in the sea otter, 414
Meienberger, C.:
(see Nagy, K. A.)
Merritt, J. F.:
Seasonal thermogenesis and changes in body mass of masked shrews, Sorex cinereus, 1020
Meserve, P. L., J. A. Yunger, J. R. Gutiérrez, L. C. Contreras, W. B. Milstead, B. K. Lang, K. L. Cramer, S. Herrera, V. O. Lagos, S. I. Silva, E. L. Tabilo, M-A. Torrealba, and F. M. Jaksic:
Heterogeneous responses of small mammals to an El Niño Southern Oscillation event in north-central semiarid Chile and the importance of ecological scale, 580
Millar, J. S.:
(see Duquette, L. S.)
(see Moses, R. A.)
Mills, J. N.:
(see Childs, J. E.)
Mills, J. N., T. L. Yates, J. E. Childs, R. R. Parmenter, T. G. Ksiazek, P. E. Rollin, and C. J. Peters:
Guidelines for working with rodents potentially infected with hantavirus, 716
Milstead, W. B.:
(see Meserve, P. L.)
Morales, J. C., and J. W. Bickham:
Molecular systematics of the genus Lasiurus (Chiroptera: Vespertilionidae) based on restriction-site maps of the mitochondrial ribosomal genes, 730
Moses, R. A., G. J. Hickling, and J. S. Millar:
Variation in sex ratios of offspring in wild bushy-tailed woodrats, 1047
Mouchaty, S., J. A. Cook, and G. F. Shields:
Phylogenetic analysis of northern hair seals based on nucleotide sequence of the mitochondrial cytochrome b gene, 1178
Mudry, M.:
(see García, M.)
Nagy, K. A., C. Meienberger, S. D. Bradshaw, and R. D. Wooller:
Field metabolic rate of a small marsupial mammal, the honey possum (Tarsipes rostratus), 862
Nilssen, K.:
(see Prestrud, P.)
O'Farrell, M. J.:
(see Scheibe, J. S.)
Olson, J. G.:
(see Gage, K. L.)
Olson, W.:
(see van Zyll de Jong, C. G.)
Ostfeld, R. S.:
(see Gage, K. L.)
Owings, D. H.:
(see Boelstorff, D. E.)
Páez, E.:
(see Lima, M.)
Painter, J., C. Krajewski, and M. Westerman:
Molecular phylogeny of the marsupial genus Planigale (Dasyuridae), 406
Parmenter, R. R.:
(see Mills, J. N.)
Peitz, D. G.:
(see Lochmiller, R. L.)
Peles, J. D., M. F. Lucas, and G. W. Barrett:
Population dynamics of agouti and albino meadow voles in high-quality, grassland habitats, 1013
Pendharkar, A. P., and S. P. Goyal:
Group size and composition of the gray goral in Simbalbara Sanctuary and Darpur Reserved Forest, India, 906
Perrin, M. R.:
(see Hughes, J. J.)
Perry, E. A., S. M. Carr, S. E. Bartlett, and W. S. Davidson:
A phylogenetic perspective on the evolution of reproductive behavior in pagophilic seals of the Northwest Atlantic as indicated by mitochondrial DNA sequences, 22
Peters, C. J.:
(see Mills, J. N.)
Pillet, J-M.:
(see Arlettaz, R.)
Ponsá, M.:
(see García, M.)
Preen, A.:
Diet of dugongs: are they omnivores?, 163
Prestrud, P., and K. Nilssen:
Growth, size, and sexual dimorphism in Arctic foxes, 522
Price, M. V.:
(see Bleich, V. C.)
Protomastro, J. J.:
(see Bolković, M. L.)
Ragen, T. J., G. A. Antonelis, and M. Kiyota:
Early migration of northern fur seal pups from St. Paul Island, Alaska, 1137
Ralls, K., and P. J. White:
Predation on San Joaquin kit foxes by larger canids, 723
Randolph, J. C., G. N. Cameron, and P. A. McClure:
Nutritional requirements for reproduction in the hispid cotton rat, Sigmodon hispidus, 1113
Rasmussen, D. T.:
(see Schwartz, G. T.)
Read, A. J.:
(see Tolley, K. A.)
Rebar, C. E.:
Ability of Dipodomys merriami and Chaetodipus intermedius to locate resource distributions, 437
Reid, F. A.:
(see Williams, S. L.)
Reynolds, H.:
(see van Zyll de Jong, C. G.)

Riddle, B. R.:
Special feature: Mammalian biogeography, 281
Molecular biogeography in the pocket mice (Perognathus and Chaetodipus) and grasshopper mice (Onychomys): the late Cenozoic development of a North American aridlands rodent guild, 283
Riedman, M. L.:
(see McShane, L. J.)
Robson, S. K.:
(see Kunz, T. H.)
Rogovin, K. A.:
(see Heske, E. J.)
Rollin, P. E.:
(see Mills, J. N.)
Rose, K. D.:
(see Heinrich, R. E.)
Rybin, S.:
(see Arlettaz, R.)
Sabat, P., F. Bozinovic, and F. Zambrano:
Role of dietary substrates on intestinal disaccharidases, digestibility, and energetics in the insectivorous mouse-opossum (Thylamys elegans), 603
Sabol, B. M., and M. K. Hudson:
Technique using thermal infrared-imaging for estimating populations of gray bats, 1242
Samson, C., and J. Huot:
Reproductive biology of female black bears in relation to body mass in early winter, 68
Scheibe, J. S., and M. J. O'Farrell:
Habitat dynamics in Peromyscus truei: eclectic females, density dependence, or reproductive constraints?, 368
Schultz, S. R., and M. K. Johnson:
Effects of birth date and body mass at birth on adult body mass of male white-tailed deer, 575
Schwartz, G. T., D. T. Rasmussen, and R. J. Smith:
Body-size diversity and community structure of fossil hyracoids, 1088
Scott, M. D.:
(see Tolley, K. A.)
Serafini, P.:
(see Cavallini, P.)
Sevick, S. H.:
(see Studier, E. H.)
Shenbrot, G. I.:
(see Heske, E. J.)
Shields, G. F.:
(see Mouchaty, S.)
Siegel, M. I.:
(see Taylor, A. B.)
Sikes, R. S.:
Costs of lactation and optimal litter size in northern grasshopper mice (Onychomys leucogaster), 348
Sillero-Zubiri, C., and D. Gottelli:
Diet and feeding behavior of Ethiopian wolves (Canis simensis), 531
Silva, S. I.:
(see Meserve, P. L.)
Skogland, T.:
(see Strand, O.)
Skupski, M. P.:
Population ecology of the western harvest mouse, Reithrodontomys megalotis: a long-term perspective, 358
Smith, J. L. D.:
(see Joshi, A. R.)
Smith, R. J.:
Harvest rates and escape speeds in two coexisting species of montane ground squirrels, 189
(see Schwartz, G. T.)
Smolen, M. J., and J. W. Bickham:
Phylogenetic implications of chromosome evolution in Geomys, 50
Soderquist, T. D.:
Otogeny of sexual dimorphism in size among polytocous mammals: tests of two carnivorous marsupials, 376
Spencer, L. M.:
Morphological correlates of dietary resource partitioning in the African Bovidae, 448
Staedler, M. M.:
(see McShane, L. J.)
Stewart, B. S., and R. L. DeLong:
Double migrations of the northern elephant seal, Mirounga angustirostris, 196
Stoddart, D. M.:
(see Xu, Z.)
Stokes, M. K.:
Selection of refuge sites of sympatric Microtus ochrogaster and Sigmodon hispidus, 83
Stott, G. G.:
(see Tarpley, R. J.)
Straley, J.:
(see Lambertsen, R.)
Strand, O., T. Skogland, and T. Kvam:
Placental scars and estimation of litter size: an experimental test in the Arctic fox, 1220
Studier, E. H., S. Sevick, D. E. Wilson, and A. P. Brooke:
Concentrations of minerals and nitrogen in milk of Carollia and other bats, 1186
Studier, E. H., and T. H. Kunz:
Accretion of nitrogen and minerals in suckling bats, Myotis velifer and Tadarida brasiliensis, 32
Sullivan, R. M.:
(see Hafner, D. J.)
Sutton, D. A.:
Problems of taxonomy and distribution in four species of chipmunks, 843
Tabilo, E. L.:
(see Meserve, P. L.)
Tarpley, R. J., G. H. Jarrell, J. C. George, J. Cubbage, and G. G. Stott:
Male pseudohermaphroditism in the bowhead whale, Balaena mysticetus, 1267
Taylor, A. B., and M. I. Siegel:
Modeling differences in biological shape in two species of Peromyscus, 828
Teferi, T., and T. B. Herman:
Epigeal movement by Sorex cinereus on Bon Portage Island, Nova Scotia, 137
Testa, J. W.:
(see Bowyer, R. T.)

Thewissen, J. G. M., and S. A. Etnier:
Adhesive devices on the thumb of vespertilionoid bats (Chiroptera), 925
Thomas, D. W.:
Hibernating bats are sensitive to nontactile human disturbance, 940
Tolley, K. A., A. J. Read, R. S. Wells, K. W. Urigan, M. D. Scott, A. B. Irvine, and A.A. Hohn:
Sexual dimorphism in wild bottlenose dolphins (Tursiops truncatus) from Sarasota, Florida, 1190
Torrealba, M-A.:
(see Meserve, P. L.)
Ulrich, N.:
(see Lambertsen, R.)
Urigan, K. W.:
(see Tolley, K. A.)
Valone, T. J., J. H. Brown, and C. L. Jacobi:
Catastrophic decline of a desert rodent, Dipodomys spectabilis: insights from a long-term study, 428
van Zyll de Jong, C. G., C. Gates, H. Reynolds, and W. Olson:
Phenotypic variation in remnant populations of North American bison, 391
Vander Wall, S. B.:
Influence of substrate water on the ability of rodents to find buried seeds, 851
Vinson, S. B.:
(see Killion, M. J.)
Wahlström, L. K.:
(see Johansson, A.)
Ward, D.:
(see Hughes, J. J.)
Warde, W. D.:
(see Davis, R. L.)
Warren, R. J.:
(see White, L. M.)
Wells, R. S.:
(see Tolley, K. A.)
Westerman, M.:
(see Painter, J.)
Whitaker, J. O., Jr.:
(see Lacki, M. J.)
White, L. M., D. A. Hosack, R. J. Warren, and R. A. Fayrer-Hosken:
Influence of mating on duration of estrus in captive white-tailed deer, 1159
White, P. J.:
(see Ralls, K.)
Williams, S. L., M. R. Willig, and F. A. Reid:
Review of the Tonatia bidens complex (Mammalia: Chiroptera), with descriptions of two new subspecies, 612
Willig, M. R.:
(see Williams, S. L.)
Willig, M. R., and R. R. Hollander:
Secondary sexual dimorphism and phylogenetic constraints in bats: a multivariate approach, 981
Wilson, D. E.:
(see Studier, E. H.)
Wilson, M. L.:
(see Krebs, J. W.)
Wolff, J. O.:
(see Manning, T.)
Woodman, N.:
Morphological variation between Pleistocene and Recent samples of Cryptotis (Insectivora: Soricidae) from the Yucatan Peninsula, Mexico, 223
Wooller, R. D.:
(see Nagy, K. A.)
Wunder, B. B.:
(see Hammond, K. A.)
Xu, Z., D. M. Stoddart, H. Ding, and J. Zhang:
Self-anointing behavior in the rice-field rat, Rattus rattoides, 1238
Yates, T. L.:
(see Dickerman, A. W.)
(see Mills, J. N.)
Yunger, J. A.:
(see Meserve, P. L.)
Zambrano, F.:
(see Sabat, P.)
Zhang, J.:
(see Xu, Z.)
Zima, J.:
(see Arlettaz, R.)

SUBJECTS
(New names in *italic* type)

Abrocoma bennetti, 580
Abrothrix longipilis, 580
Activity:
Baiomys taylori, 141
Felis concolor, 1056
Microtus oeconomus, 88
Octodon degus, 900
Paradoxurus hermaphroditus, 1205
Peromyscus leucopus, 78
Peromyscus maniculatus, 78
Rattus norvegicus, 130
Sorex cinereus, 137
Vulpes vulpes, 130
Aepyceros melampus, 450, 464
Africa:
Bovidae (33 spp), 448
Hyracoidea (19 spp), 1088
Manis tricuspis, 472
Murine rodents (6 spp), 248
Age determination:
Tadarida brasiliensis, 769
Ailuropoda melanoleuca, 339
Ailurus fulgens, 208
Akodon:
cursor, 755
molinae, 515
olivaceus, 580
Alabama:
Myotis grisescens, 1242
Alaska:
Balaena mysticetus, 1267
Callorhinus ursinus, 1137
Lutra canadensis, 1

Alberta:
Neotoma cinerea, 1047
Alcelaphus buselaphus, 450
Alces alces, 24
Aliurus fulgens, 150
Allactaga:
elater, 801
major, 801
Allometry:
Peromyscus floridanus, 828
Peromyscus gossypinus, 828
Alopex lagopus, 522, 538, 684, 723, 1220
Alouatta caraya, 755
Amblysomus:
gunningi, 960
hottentotus, 960
iris, 960
julianae, 960
Ammospermophilus:
harrisi, 296
interpres, 296
leucurus, 296, 370, 1262
nelsoni, 296
Anatomy:
Balaenopteridae, 877
Manis javanica, 472
Manis tricuspis, 472
Miacis petilus, 148
Microchiroptera (5 spp), 925
Andalgalomys olrogi, 504
Anoura geoffroyi, 981
Antechinus:
flavipes, 385
stuartii, 385
swainsonii, 385
Antidorcas marsupialis, 451, 1097
Antilocapra americana, 577
Antrozous pallidus, 484
Apodemus, 702
agrarius, 665
flavicollis, 665
mystacinus, 570
sylvaticus, 253, 570
Arctictis binturong, 207
Arctocephalus:
australis, 1249
gazella, 1249
Arctogalidia trivirgata, 207
Argentina:
Salinomys delicatus, 514
Tolypeutes matacus, 1199
Arizona:
Chaetodipus intermedius, 437
Dipodomys merriami, 437
Dipodomys spectabilis, 428
Reithrodontomys megalotis, 358
Artibeus, 670
jamaicensis, 224, 483
lituratus, 484, 981
planirostris, 981
Arvicanthus blicki, 531
Asia:
Manis javanica, 472
Atelerix:
albiventris, 1129
pruneri, 1240
Atlantic Ocean:
Balaenopteridae, 877
Eschrichtius robustus, 857
Phocinae (8 spp), 22
Australia:
Dasyurus geoffroii, 376
Dugong dugon, 163, 1226
Phascogale tapoatafa, 376
Sminthopsine marsupials (6 spp), 406
Tarsipes rostratus, 862
Baiomys taylori, 141, 363
Balaena mysticetus, 898, 1267
Balaenoptera:
acutorostrata, 877
borealis, 877
edeni, 877
musculus, 877
physalus, 877, 1268
Balantiopteryx plicata, 1129
Bandicota indica, 665
Bassaricyon gabbi, 150
Bassariscus:
astutus, 150, 698
sumichrasti, 208
Beatragus hunteri, 450
Behavior:
Canis simensis, 531
Capreolus capreolus, 123
Chaetodipus intermedius, 437
Dipodomys agilis, 646
Dipodomys merriami, 437
Dipodomys stephensi, 646
Dugong dugon, 1226
Felis concolor, 1056
Microtus oeconomus, 88
Myotis lucifugus, 940
Myotis septentrionalis, 940
Paradoxurus hermaphroditus, 1205
Phocinae (8 spp), 22
Rattus rattoides, 1238
Spermophilus beecheyi, 551
Sus scrofa, 993
Tayassu tajacu, 993
Bering Sea:
Callorhinus ursinus, 1137
Bettongia:
gaimardi, 340
lesueur, 340
penicillata, 340
Biogeography:
Cynomys (10 spp), 100
Endangered mammals, 335
Introduction, 281
Mammalian latitudinal gradient, 322
North American aridlands, 283
North American rodents (17 spp), 283
Ochotona princeps, 302
Bison bison, 449
Blarina:
brevicauda, 79, 702, 1028
carolinensis, 564
Body mass:
Hyracoidea (19 spp), 1088
Miacis petilus, 148
Odocoileus virginianus, 575

Sorex cinereus, 1020
Body size:
Spermophilus lateralis, 189
Tamias minimus, 189
Bolomys lasiurus, 755
Bos:
bison, 391
taurus, 24
Brazil:
Chiroptera (19 spp), 981
Didelphis albiventris, 750
Gracilinanus agilis, 750
Oryzomyine rodents (6 spp), 750
Breeding:
Odocoileus virginianus, 1159
Bubalus bubalis, 341
Bunohyrax:
fajumensis, 1089
major, 1089
Calcochloris obtusirostris, 960
California:
Canis latrans, 723
Enhydra lutris, 414
Felis concolor, 1056
Peromyscus truei, 368
Spermophilus beecheyi, 551
Tamias (4 spp), 843
Vulpes macrotis, 723
Vulpes vulpes, 723
Callithrix jacchus, 755
Callorhinus ursinus, 1137, 1249
Calomys:
callosus, 669, 755
laucha, 673
musculinus, 504, 669
tener, 755
Caluromys:
lanatus, 812
philander, 353, 1213
Canis:
adustus, 542
aureus, 539, 570
familiaris, 564
latrans, 528, 539, 564, 697, 723, 1004, 1060
lupus, 539, 683, 723
mesomelas, 539, 542, 640
rufus, 339
simensis, 531
Capreolus capreolus, 123
Carollia:
brevicauda, 1187
castanea, 1187
perspicillata, 483, 981, 1187
Castor canadensis, 700
Cebus:
apella, 755, 1128
capucinus, 1128
Cephalophus sylvicultor, 450
Cercocebus:
albigena, 1128
torquatus, 1128
Cercopithecus aethiops, 424
Cervus elaphus, 577
Chaetodipus:
formosus, 287
hispidus, 287
intermedius, 287, 437
penicillatus, 287, 432
pernix, 287
Chalinolobus morio, 484
Chile:
Octodon degus, 627, 900
Small mammals (7 spp), 580
Thylamys elegans, 603
Chironectes minimus, 755
Chlorotalpa:
dutieae, 960
sclateri, 960
Choeroniscus godmani, 947
Chromosomes:
Chrysochloridae (9 spp), 960
Geomys (5 spp), 50
Murine rodents (9 spp), 248
Saimiri boliviensis, 497
Chrysochloris asiatica, 960
Chrysospalax trevelyani, 960
Cladistics:
Chrysochloridae (9 spp), 960
Clethrionomys, 700
gapperi, 79
glareolus, 665, 702, 867, 1100, 1120
rufocanus, 1100
rutulis, 668
Color pattern:
Sorex cinereus, 937
Colorado:
Spermophilus lateralis, 189
Tamias minimus, 189
Communication:
Dugong dugon, 1226
Enhydra lutris, 414
Community ecology:
Didelphis albiventris, 750
Gracilinanus agilis, 750
Oryzomyine rodents (6 spp), 750
Sus scrofa, 784
Tayassu tajacu, 784
Community structure:
Hyracoidea (19 spp), 1088
Competition:
Clethrionomys glareolus, 1100
Sus scrofa, 784
Tayassu tajacu, 784
Conepatus, 224
Connochaetes taurinus, 450
Conservation biology:
Bos bison, 391
Endangered mammals, 335
Crocidura:
baileyi, 539
fumosa, 539
suaveolens, 570
Crocuta crocuta, 481, 534
Cryptotis:
mayensis, 223
mexicanus, 223
nigrescens, 223
parva, 223
Ctenomys mendocinus, 515
Cuon alpinus, 539

Cynictis penicillata, 570, 683
Cynocephalus, 1129
Cynomys:
gunnisoni, 100, 552, 697
hibbardi, 102
leucurus, 101, 559, 697
ludovicianus, 100, 697
mexicanus, 100
niobrarius, 100
parvidens, 101
vetus, 102
Cynopterus horsfieldi, 1187
Cystophora cristata, 22, 1178
Dama dama, 575
Damaliscus:
dorcas, 450
lunatus, 450
Dasyprocta, 755
Dasypus:
novemcinctus, 755
septemcinctus, 755
Dasyurus:
geoffroii, 376
viverrinus, 385
Delphinapterus leucas, 1268
Demography:
Microtus ochrogaster, 1004
Reithrodontomys megalotis, 358
Dendrolagus matschiei, 219
Density:
Didelphis albiventris, 750
Gracilinanus agilis, 750
Oryzomine rodents (6 spp), 750
Small mammals (7 spp), 580
Description:
Myosorex longicaudatus *boosmani*, 1085
Salinomys delicatus, 514
Salinomys, 505
Tonatia saurophila *bakeri*, 622
Tonatia saurophila *maresi*, 623
Desmodus rotundus, 484, 684, 986
Dicerorhinus sumatrensis, 1097
Dicrostonyx groenlandicus, 232
Didelphis:
albiventris, 750
marsupialis, 224, 809
virginiana, 224, 564, 708, 1064, 1213
Didermocercus sumatrensis, 341
Diet:
Bovidae (33 spp), 448
Canis simensis, 531
Carnivores (8 spp), 206
Dugong dugon, 163
Herpestes auropunctatus, 569
Mammals (23 spp), 1127
Myotis grisescens, 1256
Otocyon megalotis, 542
Otonycteris hemprichi, 873
Thylamys elegans, 603
Tolypeutes matacus, 1199
Digestion:
Dicrostonyx groenlandicus, 232
Microtus ochrogaster, 232
Octodon degus, 627
Thylamys elegans, 603
Dipodomys, 696
agilis, 646
californicus, 296
compactus, 296
deserti, 296, 1262
heermanni, 296
ingens, 341
merriami, 296, 432, 437, 1262
nitratoides, 296
ordii, 296, 432, 1262
panamintinus, 296, 370, 1262
spectabilis, 296, 359, 428
stephensi, 646
Disease:
Mammals (21 spp), 681
Mammals (26 spp), 665
Mammals (36 spp), 697
Distribution:
Eschrichtius robustus, 857
Myosorex longicaudatus, 1071
Tamias (4 spp), 843
Diving:
Mirounga angustirostris, 196
Dolichotis patagonum, 515
Dugong dugon, 163, 1226
Dusicyon, 515
griseus, 538
gymnocercus, 538
Dyacopterus spadiceus, 1187
Ecology:
Didelphidae (8 spp), 809
Dipodomys spectabilis, 428
Small mammals (7 spp), 580
Ectoparasites:
Mammals (36 spp), 697
Ectophylla alba, 484
Edge effect:
Mammals (14 spp), 562
Eidolon helvum, 682
Eira:
barbara, 207
eira, 150
Elaphus maximus, 341
Electrophoresis:
Ochotona princeps, 302
Oryzomyine rodents (14 spp), 172
Eligmodontia, 504
marica, 512
Energetics:
Carnivores (8 spp), 206
Onychomys leucogaster, 348
Tarsipes rostratus, 862
Thylamys elegans, 603
Enhydra lutris, 414
Eocene:
Miacis petilus, 148
Epomophorus:
gambianus, 483
wahlbergi, 484
Epomops buettikoferi, 485
Eptesicus:
furinalis, 224, 733, 984
fuscus, 482, 684, 1242
nilssoni, 486
regulus, 484

sagittula, 484
serotinus, 682
Equus grevyi, 340
Erignathus barbatus, 22, 1178
Erinaceus:
concolor, 570
europaeus, 1129, 1240
Eschrichtius robustus, 857
Ethiopia:
Canis simensis, 531
Euderma maculatum, 486
Eumops:
perotis, 1128
underwoodi, 1128
Evolution:
Cryptotis mayensis, 223
Feeding mechanics:
Balaenopteridae, 877
Felis:
concolor, 1056
pardalis, 224
rufus, 566
silvestris, 1004
weidii, 150
Fennecus zerda, 150
Finland:
Nyctereutes procyonoides, 1036
Florida:
Peromyscus floridanus, 828
Peromyscus gossypinus, 828
Tursiops truncatus, 1190
Food habits:
Bovidae (33 spp), 448
Herpestes auropunctatus, 569
Myotis grisescens, 1256
Perognathus parvus, 851
Peromyscus maniculatus, 851
Tamias amoenus, 851
Tolypeutes matacus, 1199
Foraging:
Canis simensis, 538
Chaetodipus intermedius, 437
Dipodomys merriami, 437
Dugong dugon, 163
Gerbillurus tytonis, 638
Octodon degus, 627
Otonycteris hemprichi, 873
Rattus norvegicus, 130
Spermophilus lateralis, 189
Tamias minimus, 189
Fossa fossa, 207
Fossil:
Cryptotis mayensis, 223
Cynomys (10 spp), 100
Eschrichtius robustus, 857
Hyracoidea (19 spp), 1088
Miacis petilus, 148
French Guiana:
Caluromys philander, 1213
Function:
Microchiroptera (5 spp), 925
Galagoides:
alleni, 1129
demidovii, 1129
Gazella:
granti, 451
soemmeringi, 451
thomsoni, 451
Genetta genetta, 150
Geographic variation:
Myosorex longicaudatus, 1071
Geomys:
arenarius, 64
attwateri, 50
breviceps, 50
bursarius, 43, 50
knoxjonesi, 44, 64
lutescens, 64
personatus, 50
pinetus, 51
texensis, 50
tropicalis, 65
Georgia:
Odocoileus virginianus, 1159
Gerbillurus tytonis, 638
Giraffa camelopardis, 464
Glaucomys:
sabrinus, 703
volans, 564
Glischropus tylopus, 925
Glossophaga, 477
longirostris, 947
soricina, 981
Glossotherium, 224
Gracilinanus agilis, 750
Graomys griseoflavus, 504
Great Plains:
Bos bison, 391
Cynomys (10 spp), 100
Growth:
Arctocephalus australis, 1249
Growth rates:
Alopex lagopus, 522
Caluromys philander, 1213
Clethrionomys glareolus, 867
Dasyurus geoffroii, 376
Microtus agrestis, 867
Phascogale tapoatafa, 376
Tadarida brasiliensis, 769
Gymnobelideus leadbeateri, 339
Habitat:
Clethrionomys glareolus, 1100
Dipodomys agilis, 646
Dipodomys stephensi, 646
Lutra canadensis, 1
Mammals (14 spp), 562
Microtus ochrogaster, 83
Microtus pennsylvanicus, 1013
Peromyscus truei, 368
Sigmodon hispidus, 83
Sylvilagus floridanus, 1164
Tamias striatus, 12
Halichoerus grypus, 22, 1178
Helogale:
parvula, 569
undulata, 1210
Hemiechinus auritus, 1240
Herpestes auropunctatus, 569, 684, 1206
Heterocephalus glaber, 481

Heteromys gaumeri, 224
Hibernation:
Myotis lucifugus, 940
Myotis septentrionalis, 940
Hipposideros:
caffer, 1128
commersoni, 1128
speoris, 485
Hippotragus:
equinus, 450
niger, 450
Holochilus brasiliensis, 174
Home range:
Lutra canadensis, 1
Sigmodon hispidus, 819
Spermophilus beecheyi, 551
Stylodipus telum, 800
Sus scrofa, 993
Tamias striatus, 12
Tayassu tajacu, 993
Homo sapiens, 1129
Hyaena brunnea, 481
Hybridization:
Geomys bursarius, 43
Geomys knoxjonesi, 44
Hydrochaeris hydrochaeris, 755
Hypsignathus monstrosus, 483
Ichneumia albicauda, 1210
Illinois:
Carnivora (8 spp), 1004
Mammals (14 spp), 562
Sylvilagus floridanus, 1010
Immunity:
Sigmodon hispidus, 912
India:
Nemorhaedus goral, 906
Kansas:
Microtus ochrogaster, 83
Onychomys leucogaster, 348
Sigmodon hispidus, 83
Kentucky:
Myotis grisescens, 1256
Kirghizstan:
Otonycteris hemprichi, 873
Kobus:
ellipsiprymnus, 450
kob, 450
leche, 450
vardoni, 450
Korcula Island:
Herpestes auropunctatus, 569
Kvabebihyrax kachethicus, 1089
Lactation:
Caluromys philander, 1213
Myotis velifer, 32
Onychomys leucogaster, 348
Tadarida brasiliensis, 32
Lagorchestes hirsutus, 340
Lagostrophus fasciatus, 340
Lasionycteris noctivagans, 686
Lasiorhinus krefftii, 340
Lasiurus:
blossevilli, 730
borealis, 684, 730
cinereus, 491, 684, 730
ega, 730
insularis, 730
intermedius, 684, 730
pfeifferi, 730
seminolus, 684, 730
xanthinus, 730
Learning:
Chaetodipus intermedius, 437
Dipodomys merriami, 437
Lemmiscus curtatus, 108
Lemmus:
lemmus, 1221
sibiricus, 1120
Leontopithecus rosalia, 341
Leporillus conditor, 340
Leptonychotes weddelli, 24, 1180
Leptonycteris curasoae, 947
Lepus:
americanus, 700
californicus, 700
starcki, 531
Litocranius walleri, 453
Litter size:
Alopex lagopus, 1220
Onychomys leucogaster, 348
Ursus americanus, 68
Lonchophylla mordax, 985
Lophuromys:
flavopunctatus, 532
melanonyx, 531
Louisiana:
Odocoileus virginianus, 575
Lutra canadensis, 1
Lycaon pictus, 539
Macaca fasicularis, 1129
Macropus giganteus, 608
Macrotis lagotis, 340
Macrotus californicus, 490
Malaysia:
Pteropid bats (3 spp), 1187
Manis:
dalmanni, 477
gigantea, 477
javanica, 472
pentadactyla, 477
temminckii, 477
tricuspis, 472
Marking:
Capreolus capreolus, 123
Marmosa:
canescens, 224
murina, 812
pallidior, 515
Marmosops noctivagus, 809
Marmota:
flaviventris, 552
monax, 552, 566, 687
Martes:
americana, 24, 150, 698
foina, 570
pennanti, 150
Maryland:
Peromyscus leucopus, 78
Peromyscus maniculatus, 78
Mastomys, 669

coucha, 248
erythroleucus, 248
huberti, 248
natalensis, 248, 675
Mating system:
Tursiops truncatus, 1190
Mazama americana, 755
Megaderma lyra, 485
Megaerops ecaudatus, 1187
Megalohyrax:
championi, 1089
eocaenus, 1089
Megaptera novaeangliae, 877
Meles meles, 481
Mephitis, 224, 1067
mephitis, 150, 564, 683, 698
Meriones meridianus, 801
Metabolism:
Carnivores (8 spp), 206
Choeroniscus godmani, 947
Glossophaga longirostris, 947
Leptonycteris curasoae, 947
Sorex cinereus, 1020
Tarsipes rostratus, 862
Metachirus nudicaudatus, 812
Miacis:
deutschi, 148
exiguus, 148
kessleri, 149
parvivorus, 149
petilus, 148
winkleri, 148
Micoureus cinereus, 812
Microhyrax lavocati, 1089
Microryzomys minutus, 174
Microsyops, 1129
Microtus, 700
agrestis, 867, 1100
californicus, 666, 697, 716
canicaudus, 1149
fortis, 709
miurus, 1100
montanus, 1100
ochrogaster, 83, 232, 1004, 1221
oeconomus, 88
pennsylvanicus, 665, 716, 1013, 1100
pinetorum, 564, 708, 1120
Migration:
Callorhinus ursinus, 1137
Mirounga angustirostris, 196
Milk:
Bats (8 spp), 1186
Myotis velifer, 32
Tadarida brasiliensis, 32
Mimon:
bennetti, 224
cozumelae, 224
Miniopterus australis, 484
Mirounga angustirostris, 196, 424
Mitochondrial DNA:
Lasiurus (9 spp), 730
North American rodents (17 spp), 283
Phocinae (8 spp), 22, 1178
Sminthopsine marsupials (6 spp), 406
Molecular genetics:
Geomys bursarius, 44
Geomys knoxjonesi, 44
Molossus molossus, 981
Monachus schauinslandi, 24, 1180
Monodelphis:
americana, 755
domestica, 1218
Monodon monoceros, 1190
Monophyllus, 477
Mormoops megalophylla, 224, 490
Morphology:
Bovidae (33 spp), 448
Dicrostonyx groenlandicus, 232
Microtus ochrogaster, 232
Otocyon megalotis, 542
Peromyscus floridanus, 828
Peromyscus gossypinus, 828
Morphometrics:
Chiroptera (19 spp), 981
Tamias (4 spp), 843
Tursiops truncatus, 1190
Mungos mungo, 150, 569
Mus:
domesticus, 570
musculus, 145, 224, 253, 349, 596, 669, 1221
Mustela:
frenata, 224, 564, 698, 1004
lutreola, 341
nigripes, 341
nivalis, 1004
sibirica, 1238
vison, 150, 1004
Myology:
Manis javanica, 472
Manis tricuspis, 472
Myomys daltoni, 248
Myosorex:
cafer, 1071
geata, 1084
longicaudatus, 1071
schalleri, 1084
varius, 1071
Myotis:
adversus, 492
bechsteini, 491
emarginatus, 491
grisescens, 482, 1242, 1256
lucifugus, 482, 940, 1128, 1258
myotis, 486
riparia, 981
septentrionalis, 940
sodalis, 485
velifer, 32, 1128
Myoxus glis, 570
Myrmecobius fasciatus, 340
Myrmecophaga, 477
Myzopoda aurita, 925
Namibia:
Gerbillurus tytonis, 638
Nandinia binotata, 150, 207
Nasua:
narica, 208
nasua, 208
Neacomys spinosus, 174

Nectomys squamipes, 174, 750
Nemorhaedus goral, 906
Neofelis nebulosa, 1210
Neomys:
anomalus, 1028
fodiens, 1028
Neoplatymops mattogrossensis, 985
Neotoma, 697
albigula, 432
cinerea, 703, 1047
fuscipes, 702
lepida, 370
Nepal:
Paradoxurus hermaphroditus, 1205
Nevada:
Rodents (6 spp), 1260
New Mexico:
Chaetodipus intermedius, 437
Dipodomys merriami, 437
Noctilio:
albiventris, 485
leporinus, 485, 985
North America:
Ochotona princeps, 302
Rodents (17 spp), 283
Norway:
Alopex lagopus, 522
Microtus oeconomus, 88
Notomys fuscus, 340
Nova Scotia:
Sorex cinereus, 137
Nutrition:
Bats (8 spp), 1186
Myotis velifer, 32
Octodon degus, 627
Sigmodon hispidus, 912, 1113
Sylvilagus floridanus, 1164
Tadarida brasiliensis, 32
Nyctalus:
lasiopterus, 482
noctula, 481
Nyctereutes procyonoides, 683, 1036
Nycteris grandis, 485
Nycticeius humeralis, 482
Nyctimene robinsoni, 484
Nyctophilus:
geoffroyi, 485
gouldi, 485
Ochotona collaris, 302
Ochrotomys nuttalli, 564
Octodon:
degus, 580, 627, 900
lunatus, 582
Odocoileus:
hemionus, 24, 1061
virginianus, 24, 349, 566, 575, 702, 1159
Oecomys:
bicolor, 173, 750
concolor, 750
Ohio:
Microtus pennsylvanicus, 1013
Oil spill:
Lutra canadensis, 1
Oklahoma:
Sylvilagus floridanus, 1164
Oligoryzomys:
chacoensis, 173
delicatus, 173
destructor, 173
eliurus, 183, 750
flavescens, 173
fornesi, 173
fulvescens, 173, 224
longicaudatus, 173, 580
microtis, 173
nigripes, 173
Ondatra zibethicus, 700
Onychomys, 432
arenicola, 287
bensoni, 292
leucogaster, 287, 348, 697
torridus, 287
Orcinus orca, 424
Oregon:
Microtus canicaudus, 1149
Tamias senex, 843
Oreotragus oreotragus, 450
Oryx gazella, 450
Oryzomys, 699
capito, 173, 750
couesi, 224
keaysi, 174
nitidus, 174
palustris, 145, 173
subflavus, 174
Otocyon megalotis, 542
Otomys typus, 531
Otonycteris hemprichii, 873
Otonyctomys hatti, 224
Ototylomys phyllotis, 224, 224
Ourebia ourebi, 450
Pachyhyrax crassidentatus, 1089
Pacific Ocean:
Balaenopteridae, 877
Callorhinus ursinus, 1137
Mirounga angustirostris, 196
Paleoecology:
North American rodents (17 spp), 283
Paleontology:
Cynomys (10 spp), 100
Hyracoidea (19 spp), 1088
Panthera:
pardus, 1210
tigris, 341, 1206
Papio cynocephalus, 481
Paradoxurus hermaphroditus, 150, 207, 1205
Parantechinus apicalis, 340
Parapliohyrax ngororaensis, 1089
Pelage:
Microtus pennsylvanicus, 1013
Pennsylvania:
Sorex cinereus, 1020
Peripheral populations:
Endangered mammals, 335
Perognathus:
amplus, 289
apache, 289
fasciatus, 287

flavus, 287, 428
inornatus, 288
longimembris, 287, 1262
merriami, 289
parvus, 287, 370, 851
Peromyscus:
boylii, 667
eremicus, 296, 432
floridanus, 828
gossypinus, 828
leucopus, 78, 145, 349, 564, 702, 1120
maniculatus, 78, 349, 368, 432, 664, 697, 702, 716, 851
mexicanus, 596
truei, 368, 667
yucatanicus, 224
Peropteryx macrotis, 985
Peru:
Didelphidae (8 spp), 809
Phyllostomid bats (4 spp), 1187
Phacochoerus aethiopicus, 1097
Phascogale:
calura, 340
tapoatafa, 376
Phascolarctos cinereus, 608
Phenotypic variation:
Bos bison, 391
Philander andersoni, 809
Phoca:
fasciata, 1178
groenlandica, 22, 1178
hispida, 22, 1178
largha, 1178
vitulina, 22, 1178
Phocoena phocoena, 1190
Phyllostomus:
discolor, 981
hastatus, 481, 985, 1187
Phyllotis:
darwini, 580
gerbillus, 518
Phylogeny:
Geomys (5 spp), 50
Lasiurus (9 spp), 730
Murine rodents (9 spp), 248
Oryzomyine rodents (14 spp), 172
Phocinae (8 spp), 22, 1178
Sminthopsine marsupials (6 spp), 406
Physeter catodon, 1190
Pipistrellus:
nanus, 483, 733, 925
pipistrellus, 1258
Planigale:
gilesi, 385, 406
ingrami, 406
maculata, 406
novaeguinea, 406
tenuirostris, 406
Pleistocene:
Cryptotis mayensis, 223
Cynomys (6 spp), 102
Pliocene:
Cynomys hibbardi, 102
Cynomys vetus, 102
Pongo pygmaeus, 220
Population dynamics:
Clethrionomys glareolus, 1100
Dipodomys spectabilis, 428
Microtus pennsylvanicus, 1013
Small mammals (7 spp), 580
Sus scrofa, 993
Tayassu tajacu, 993
Population ecology:
Reithrodontomys megalotis, 358
Population genetics:
Ochotona princeps, 302
Population size:
Myotis grisescens, 1242
Postschizotherium chardini, 1089
Potos flavus, 150, 207
Praomys tullbergi, 248
Predation:
Canis latrans, 723
Carnivora (8 spp), 1004
Gerbillurus tytonis, 638
Microtus ochrogaster, 1004
Rattus norvegicus, 130
Spermophilus lateralis, 189
Tamias minimus, 189
Vulpes macrotis, 723
Vulpes vulpes, 130
Procavia capensis, 534
Procyon lotor, 208, 224, 481, 564, 683, 698, 702, 1004
Proechimys longicaudatus, 755
Proteles cristatus, 215
Pseudohermaphroditism:
Balaena mysticetus, 1267
Pseudomys:
fumeus, 340
oralis, 340
shortridgei, 340
Pteronotus:
parnellii, 224
personatus, 608
Pteronura brasiliensis, 424
Pteropus:
giganteus, 220
hypomelanus, 220
poliocephalus, 483
Putorius putorius, 1240
Pygeretmus pumilio, 801
Quebec:
Myotis lucifugus, 940
Myotis septentrionalis, 940
Ursus americanus, 68
Radiotelemetry:
Callorhinus ursinus, 1137
Felis concolor, 1056
Paradoxurus hermaphroditus, 1205
Sigmodon hispidus, 819
Raphicerus campestris, 464
Rattus:
norvegicus, 130, 570, 666, 697, 716
rattoides, 1238
rattus, 253, 570, 667, 695, 755
Redunca:
arundinum, 450
fulvorufula, 450
redunca, 450, 538

Refuge site:
Microtus ochrogaster, 83
Sigmodon hispidus, 83
Reithrodontomys:
fulvescens, 145
gracilis, 224
megalotis, 358, 432, 666, 716
Reproduction:
Arctocephalus australis, 1249
Didelphidae (8 spp), 809
Nyctereutes procyonoides, 1036
Odocoileus virginianus, 1159
Peromyscus mexicanus, 596
Peromyscus truei, 368
Sigmodon hispidus, 1113
Ursus americanus, 68
Rhabdomys pumilio, 640
Rhinoceros:
sondaicus, 339
unicornis, 339
Rhinolophus:
hildebrandti, 486
simulator, 1129
Rhinopoma hardwickei, 1129
Rhipidomys mastacalis, 750
Rhynchonycteris naso, 484
Roosts:
Chiroptera (43 spp), 481
Rousettus amplexicaudatus, 1129
Russia:
Stylodipus telum, 800
Saccopteryx:
bilineata, 484
leptura, 484
Safety techniques:
Peromyscus maniculatus, 716
Rattus norvegicus, 716
Sigmodon hispidus, 716
Saghatherium:
bowni, 1089
humarum, 1089
Saimiri:
boliviensis, 497
sciureus, 498
Salinomys delicatus, 514
Sciurus:
carolinensis, 702
yucatanensis, 224
Scotophilus leucogaster, 485
Seed Caching:
Rodents (6 spp), 1260
Selenohyrax chatrathi, 1089
Sex ratio:
Neotoma cinerea, 1048
Ursus americanus, 68
Sexual dimorphism:
Alopex lagopus, 522
Chiroptera (19 spp), 981
Dasyurus geoffroii, 376
Phascogale tapoatafa, 376
Tursiops truncatus, 1190
Sigmodon, 363
alstoni, 669
hispidus, 83, 141, 224, 353, 665, 708, 716, 819, 912, 1113
Size:
Clethrionomys glareolus, 867
Microtus agrestis, 867
Tadarida brasiliensis, 769
Sminthopsis:
crassicaudata, 407
longicaudata, 340
murina, 385, 407
Social structure:
Nemorhaedus goral, 906
Spermophilus beecheyi, 551
Sogdohyrax soricus, 1089
Sorex:
araneus, 1028
cinereus, 79, 137, 937, 1020
dispar, 79
fumeus, 79
minutus, 1028
vagrans, 1029
South Africa:
Chrysochloridae (9 spp), 960
Myosorex longicaudatus, 1071
Otocyon megalotis, 542
Steatomys pratensis, 241
South America:
Oryzomyine rodents (14 spp), 172
Spatial association:
Nemorhaedus goral, 906
Stylodipus telum, 800
Species richness:
Western hemisphere mammals, 322
Spermophilus:
armatus, 697
beecheyi, 551, 697
beldingi, 193, 697
columbianus, 551, 703
elegans, 697
lateralis, 189, 697
richardsonii, 551
saturatus, 193, 349
tereticaudus, 551
townsendii, 697
tridecemlineatus, 552
variegatus, 551, 697
Spilogale:
angustifrons, 224
gracilis, 698
Steatomys pratensis, 240
Stenella coeruleoalba, 1268
Stenocephalemys:
albocaudata, 532
griseicauda, 532
Stylodipus telum, 800
Suncus:
etruscus, 570
murinus, 665, 1240
Suricata suricatta, 569
Survival:
Myotis lucifugus, 940
Myotis septentrionalis, 940
Sus scrofa, 784, 993
Sweden:
Capreolus capreolus, 123
Clethrionomys glareolus, 867, 1100
Microtus agrestis, 867

Sylvicapra grimmia, 450
Sylvilagus, 224
auduboni, 700
brasiliensis, 755
floridanus, 700, 1010, 1164
Syncerus caffer, 449
Systematics:
Lasiurus (9 spp), 730
Oryzomyine rodents (14 spp), 172
Phyllotine rodents (5 spp), 504
Tonatia bidens, 612
Tonatia saurophila, 612
Tachyglossus aculeatus, 477
Tachyoryctes macrocephalus, 531
Tadarida:
brasiliensis, 32, 686, 769, 1186
midas, 485
pumila, 485
Tamandua, 477
Tamias, 667
amoenus, 443, 703, 843, 851
merriami, 370
minimus, 189
quadrimaculatus, 843
senex, 843
sibiricus, 1238
speciosus, 843
striatus, 12, 79, 564, 702
Tamiasciurus hudsonicus, 79, 107, 703
Taphozous mauritianus, 1129
Tapirus indicus, 1097
Tarsipes rostratus, 862
Taurotragus oryx, 450
Taxidea taxus, 698
Taxonomy:
Chrysochloridae (9 spp), 960
Myosorex longicaudatus, 1071
Tamias (4 spp), 843
Tonatia bidens, 612
Tonatia saurophila, 612
Tayassu tajacu, 224, 784, 993
Technique:
Estimation of litter size, 1220
Fluorescent pigments, 1260
Population-size estimators, 1149
Thermal infared-imaging, 1242
Tertiary:
Hyracoidea (19 spp), 1088
Texas:
Baiomys taylori, 141
Geomys bursarius, 43
Geomys knoxjonesi, 44
Myotis velifer, 32
Sigmodon hispidus, 819, 1113
Sus scrofa, 784, 993
Tadarida brasiliensis, 32, 769
Tayassu tajacu, 784, 993
Thermogenesis:
Sorex cinereus, 1020
Thermoregulation:
Octodon degus, 900
Steatomys pratensis, 240
Thylamys elegans, 580, 603
Thyrohyrax:
domorictus, 1089
litholagus, 1089
meyeri, 1089
Thyroptera tricolor, 483, 925
Titanohyrax:
angustidens, 1089
mongereaui, 1089
ultimus, 1089
Tolypeutes matacus, 1199
Tonatia:
bidens, 612, 985
brasiliensis, 619
carrikeri, 619
evotis, 612
minuta, 619
nicaraguae, 619
saurophila, 612, 983
schulzi, 619
silvicola, 619
venezuelae, 619
Tooth enamel:
Mammals (23 spp), 1127
Torpor:
Steatomys pratensis, 240
Tarsipes rostratus, 862
Trachops cirrhosus, 984
Tragelaphus:
buxtoni, 534
euryceros, 450
imberbis, 450
scriptus, 450
spekei, 450
strepiceros, 450
Trichechus:
inunguis, 1234
manatus, 1234
Tursiops truncatus, 1190
Tylomys nudicaudatus, 224
Tylonycteris:
pachypus, 485, 925
robustula, 485
United Kingdom:
Rattus norvegicus, 130
Vulpes vulpes, 130
Urocyon cinereoargenteus, 224, 564, 684, 698, 1004
Uroderma bilobatum, 483
Ursus:
americanus, 68, 698
ursinus, 208
Uruguay:
Arctocephalus australis, 1249
Vampyrodes caraccioli, 484
Vampyrops lineatus, 981
Vampyrum spectrum, 485
Venezuela:
Choeroniscus godmani, 947
Glossophaga longirostris, 947
Leptonycteris curasoae, 947
Veracruz:
Peromyscus mexicanus, 597
Virginia:
Tamias striatus, 12
Viverra zibetha, 1205
Vocalization:
Dugong dugon, 1226
Enhydra lutris, 414

Vulpes, 698
chama, 542
ferrilata, 539
fulva, 1004
macrotis, 723
vulpes, 130, 528, 538, 564, 683, 723, 1238
Wyoming:
Miacis petilus, 148
Xerus inauris, 481
Yucatán:
Cryptotis mayensis, 223
Zalophus californianus, 424
Zoonoses:
Mammals (21 spp), 681
Mammals (26 spp), 665
Mammals (36 spp), 697
Zygodontomys brevicaudata, 674
Zygogeomys trichopus, 313

Reviews

Choate, J. R., J. K. Jones, Jr., and C. Jones:
Handbook of mammals of the south-central states, reviewed by D. M. Armstrong, 1282
Findley, J. S.:
Bats: a community perspective, reviewed by R. C. B. Utzurrum and T. H. Kunz, 264
Jefferson, T. A., S. Leatherwood, and M. A. Webber:
Marine mammals of the world, reviewed by H. D. Walley, 975
Meffe, G. K., C. R. Carroll, and contributors:
Principles of conservation biology, reviewed by J. L. Morrison, 972
Riedman, M. L. and J. A. Estes:
The sea otter (Enhydra lutris): behavior, ecology, and natural history, reviewed by J. J. Flynn, 263
Stangl, F. B., Jr., W. W. Dalquest, and R. R. Hollander:
Evolution of a desert mammalian fauna: a 10,000-year history of mammals from Culberson and Jeff Davis counties, Trans-Pecos Texas, reviewed by A. H. Harris, 1290
Szalay, F.S.:
Evolutionary history of the marsupials and an analysis of osteological characters, reviewed by J. A. W. Kirsch, 1283
Wainwright, P. C., and S. M. Reilly (eds.):
Ecological morphology: integrative organismal biology, reviewed by W. L. Gannon, 1288

Letters

Hauber, D. J.:
The International Code of Zoological Nomenclature as a guide to the pronunciation of zoological Latin, 652

Obituary

Francis Hollis Fay (1927-1994):
Appreciation by Robert L. Rausch, 1276